

Aan de staatssecretaris van  
Infrastructuur en Milieu  
Mevrouw W.J. Mansveld  
Postbus 30945  
2500 GX Den Haag

**DATUM** 07 november 2013  
**KENMERK** CGM/131107-01  
**ONDERWERP** Advies: Import and processing of insect resistant and herbicide tolerant cotton  
281-24-236x3006-210-23xMON88913

Geachte mevrouw Mansveld,

Naar aanleiding van de adviesvraag betreffende het dossier EFSA/GMO/NL/2009/68 voor import en verwerking van de genetisch gemodificeerde katoenlijn 281-24-236x3006-210-23xMON88913, ingediend door Agrigenetics, Inc. d/b/a Mycogen Seeds, deelt de COGEM u het volgende mee.

**Samenvatting:**

De COGEM is gevraagd te adviseren over de import en de verwerking van de genetisch gemodificeerde katoenlijn 281-24-236x3006-210-23xMON88913. Deze katoenlijn brengt de genen *cry1F*, *cry1Ac*, *pat* en *cp4 epsps* tot expressie waardoor het gewas resistent is tegen bepaalde insecten uit de orde Lepidoptera en tolerant voor glufosinaat-ammonium en glyfosaat bevattende herbiciden.

In Europa komen geen wilde verwanten van katoen voor. De katoenplant is sterk koudegevoelig en heeft hoge temperaturen en veel water nodig om te kiemen en te groeien. Katoen kan niet overleven in het Noordwest-Europese klimaat. De ingebrachte eigenschappen zorgen er niet voor dat 281-24-236x3006-210-23xMON88913 in Noordwest-Europa zou kunnen groeien. Daarom acht de COGEM de kans verwaarloosbaar klein dat het incidenteel morsen van de katoenzaden leidt tot verspreiding of vestiging van de genetisch gemodificeerde katoenlijn 281-24-236x3006-210-23xMON88913 in Noordwest-Europa.

Omdat andere instanties een voedselveiligheidsbeoordeling uitvoeren heeft de COGEM bij deze vergunningaanvraag de risico's van incidentele consumptie niet beoordeeld. Concluderend acht de COGEM de risico's voor mens en milieu verwaarloosbaar klein bij import en verwerking van gg-katoen 281-24-236x3006-210-23xMON88913.



# **Import and processing of insect resistant and herbicide tolerant cotton 281-24-236x3006-210-23xMON88913**

## **COGEM advisory report CGM/131107-01**

### **Summary**

*The present application (EFSA/GMO/NL/2009/68) concerns import and processing for use in feed and food of the genetically modified cotton event 281-24-236x3006-210-23xMON88913. Cultivation is not part of this application.*

*Cotton line 281-24-236x3006-210-23xMON88913 expresses the Cry1F, Cry1Ac PAT and CP4 EPSPS proteins, conferring resistance to certain lepidopteran insect pests and tolerance to glufosinate-ammonium and glyphosate containing herbicides.*

*281-24-236x3006-210-23xMON88913 was produced by means of conventional breeding of two genetically modified parental cotton lines. Previously, COGEM issued positive opinions on the import and processing of the parental lines 281-24-236x3006-210-23 and MON88913.*

*In Northwestern Europe, no wild relatives of cotton are present. Modern cotton cultivars do not possess any of the attributes commonly associated with problematic weeds. Cotton cannot survive in Northwestern Europe due to the climate. The introduced traits will not introduce a potential to establish feral populations. Therefore, COGEM is of the opinion that incidental spillage of 281-24-236x3006-210-23xMON88913 seeds will not pose a risk to the environment in Northwestern Europe.*

*In view of the above, COGEM is of the opinion that import and processing of cotton line 281-24-236x3006-210-23xMON88913 poses a negligible risk to man and the environment. A food/feed safety assessment is carried out by other organisations. Therefore, COGEM abstains from advice on the potential risks of incidental consumption.*

### **Introduction**

The scope of the present application (EFSA/GMO/NL/2009/68) submitted by Agrigenetics, Inc. d/b/a Mycogen Seeds, concerns import and processing of cotton line 281-24-236x3006-210-23xMON88913. This line was produced by conventional crossbreeding of the genetically modified parental cotton lines 281-24-236x3006-210-23 and MON88913. It expresses the Cry1F and Cry1Ac proteins conferring resistance to certain lepidopteran insect pests. It also expresses the PAT and CP4 EPSPS proteins, conferring tolerance to glufosinate-ammonium and glyphosate herbicides. The applicant states that glufosinate-ammonium will only be used during the selection stages of the genetic modification.

### **Previous COGEM advisory reports**

In 2004, COGEM issued a positive opinion on the import and processing of parental line 281-24-236x3006-210-23.<sup>1,2</sup> In 2007, COGEM also issued a positive opinion on the import and processing of parental line MON88913.<sup>3</sup>

### Aspects of the crop

Cotton is a member of the genus *Gossypium* and belongs to the *Malvaceae* family. The majority of cultivated cotton is *Gossypium hirsutum* (90%) and *Gossypium barbadense* (5%), while *Gossypium arboreum* and *Gossypium herbaceum* are cultivated as well.<sup>4,5,6</sup> The only cultivated cotton species in Europe is *G. hirsutum*, which is grown in Greece, Spain and Bulgaria.<sup>7</sup>

Cotton plants reproduce sexually.<sup>6</sup> Cotton is predominantly a self-pollinating species, but cross-pollination may occur. Dissemination of pollen by wind is (almost) absent.<sup>5,6</sup> The pollen of cotton is large, heavy and somewhat sticky.<sup>5,6</sup> Outcrossing rates for cotton are strongly influenced by the presence of insects. Pollinators of cotton flowers include bumblebees (*Bombus* spp.), honeybees (*Apis* spp.) and other bee species (*Anthophora* spp., *Melissodes* spp. and *Halictus* spp.).<sup>5,8</sup> Hybridisation between *G. hirsutum* and *G. barbadense* may occur and can lead to viable progeny (F1). Hybrid F2 progeny contain either depauperate types or plants that closely resemble one of the parents.<sup>8</sup> Wild relatives of cotton (*Gossypium* spp.) do not occur in Northwestern Europe. Therefore, hybridisation with wild relatives cannot occur in Northwestern Europe.<sup>5</sup>

Cotton is highly sensitive to temperature and susceptible to frost. Temperature is the main factor that determines the geographic range in which cotton can be grown. Plant development ceases below a temperature of 12 °C and delays when the temperature rises above 38 °C.<sup>5,6</sup> *G. hirsutum* needs a period of 180 to 200 frost-free days for normal maturation, with an average of 150 days of suitable temperatures (averaging 21-22 °C). The optimal temperature for growth is between 30 and 35 °C.<sup>9</sup> *G. barbadense* has a longer growing season and needs 200-250 frost-free days and a lower optimum temperature for growth (25-30°C).<sup>5,10</sup> In places where cotton is cultivated as a rain-fed crop the average rainfall is 800-1200 mm.<sup>4</sup> In areas where the rainfall is less than 500 mm a year, irrigation is necessary.<sup>5</sup>

Cottonseed can be dispersed by wind, water, during transport or when feeding cattle.<sup>6</sup> In addition, cottonseed can be transported by birds or rodents. Seeds from cotton cultivars do not possess dormancy and will germinate directly if conditions are favourable.<sup>5,6</sup> Seeds usually do not survive in humid soil.<sup>6</sup> In regions with mild and dry winters, cottonseeds may overwinter and germinate in spring. Seedlings are sensitive to competition from weeds.<sup>5</sup>

Modern cotton cultivars do not possess any of the attributes commonly associated with problematic weeds, such as dormancy, persistence in seed banks, germination under adverse environmental conditions, rapid vegetative growth, a short life cycle, very high seed output, high seed dispersal and long-distance dispersal of seeds. Cotton volunteers occur in cotton growing areas and may occur when cottonseed is used as livestock feed. The presence of volunteer cotton is limited by soil moisture content and frost.<sup>6</sup> There are reports that *G. hirsutum* and *G. herbaceum* cotton are naturalised in some Southern European countries, e.g. Greece and Spain.<sup>11,12</sup>

### **Molecular characterization**

281-24-236x3006-210-23xMON88913 cotton is the result of a conventional breeding cross between the two genetically modified cotton lines 281-24-236x3006-210-23 and MON88913. COGEM evaluated the molecular characterization of the genetically modified parental lines in previous applications concerning import and processing and concluded that the molecular characterization of the individual parental lines is adequate.<sup>1,2,3</sup> The bioinformatics analyses of cotton line 281-24-236x3006-210-23xMON88913 were performed using up-to-date databases.

### ***Properties of the introduced genes conferring insect resistance***

Cotton line 281-24-236x3006-210-23xMON88913 contains the *cryIF* and *cryIAC* genes. These *cry* genes encode delta-endotoxins ( $\delta$ -endotoxin), which target lepidopteran insects such as the Tobacco Budworm (*Heliothis virescens*) and Pink Bollworm (*Pectinophora gossypiella* Sanders).  $\delta$ -endotoxins are solubilised in the midgut of susceptible insects and are activated by midgut proteases to release a toxin fragment. This toxin fragment binds to specific receptors on the epithelial surface of the midgut, which causes pores to open. This leads to disruption of the movement of solutes across the gut epithelium and allows gut bacteria to escape the midgut and enter the hemolymph where they cause septicaemia and death.<sup>13,14</sup>

### ***Properties of the introduced genes conferring herbicide tolerance***

281-24-236x3006-210-23xMON88913 expresses the phosphinothricin acetyltransferase (PAT) and CP4 EPSPS proteins. The *cp4 epsps* gene encodes the CP4 EPSPS protein. EPSPS is an enzyme involved in the biosynthesis of aromatic amino acids. Glyphosate inhibits EPSPS, resulting in a lack of amino acids essential for growth and development of plants. In contrast to EPSPS, the CP4 EPSPS protein is not inhibited by glyphosate and therefore the plant is tolerant to glyphosate containing herbicides.<sup>15</sup>

PAT confers tolerance to glufosinate-ammonium containing herbicides.<sup>16</sup> The active ingredient in glufosinate-ammonium is L-phosphinothricin (L-PPT), which binds to glutamine synthetase in plants. The detoxification of ammonia is thereby prevented, leading to plant death. The PAT enzyme catalyses the conversion of L-PPT to an inactive form, which does not bind glutamine synthetase.<sup>17</sup> The applicant states that glufosinate-ammonium will only be used during the selection stages of the genetic modification.

### **Environmental risk assessment**

Cotton is predominantly a self-pollinating species, but cross-pollination may occur. Wild relatives of cotton (*G. hirsutum* and *G. barbadense*) are not present in Northwestern Europe and therefore, hybridisation with wild relatives is excluded.<sup>5</sup>

Cotton plants are susceptible to frost. A reasonably high temperature (optimally 25-30 °C for *G. barbadense* and 30-35 °C for *G. hirsutum*) is required in all stages of development. For normal maturation, a period of 200 to 250 frost-free days is needed for *G. barbadense* and 180 to 200 days

for *G. hirsutum*. In addition, this period needs to have on average 150 days of suitable temperatures, averaging 21-22 °C. In the Netherlands, the summer months of May, June, July, August and September (in total 153 days) have average monthly temperatures above 12 °C, but below 18 °C.<sup>18</sup> In addition, in areas where rainfall is less than 500 mm a year, irrigation should be applied for cotton growth. In the months of May through September in the Netherlands, the average monthly precipitation does not exceed 100 mm.<sup>18</sup> Based on the above, the current Dutch climate is unsuited for cotton growth.

There is no indication that the introduced traits, which confer resistance to certain lepidopteran insects and tolerance to glufosinate-ammonium and glyphosate containing herbicides, will introduce an ability of cotton to survive in the environment. The applicant carried out an agronomic assessment for 281-24-236x3006-210-23xMON88913. The agronomic assessment does not give any indication to assume that 281-24-236x3006-210-23xMON88913 has an increased fitness compared to conventional varieties.

In view of the above, there are no reasons to assume that 281-24-236x3006-210-23xMON88913 has an increased potential for the establishment of feral populations in case of incidental spillage of cottonseed because the climate in Northwestern Europe is not suited for cotton growth.

Since 2008 COGEM abstains from giving advice on the potential risks of incidental consumption in case a food/feed assessment is already carried out by other organisations.<sup>19</sup> This application is submitted under Regulation (EC) 1829/2003, therefore a food/feed assessment is carried out by EFSA. Other organisations who advise the competent authorities can perform an additional assessment on food safety although this is not obligatory. In the Netherlands a food and/or feed assessment for Regulation (EC) 1829/2003 applications is carried out by RIKILT. Regarding the risks for food and feed, the outcome of the assessment by other organisations (EFSA, RIKILT) was not known at the moment of the completion of this advice.

### **General surveillance plan**

General surveillance (GS) has been introduced to be able to observe unexpected adverse effects of genetically modified (GM) crops on the environment. The applicant's general surveillance plan is in line with the criteria COGEM formulated in 2010.<sup>20</sup>

COGEM notes that the GS plan of cotton line 281-24-236x3006-210-23xMON88913 could be improved by a statement that raw data and analysis of monitoring data will be made available to the Competent Authorities and the European Commission.<sup>21</sup> A similar remark in general has been made by EFSA in its guidance document on the Post-Market Environmental Monitoring of genetically modified plants.<sup>22</sup>

## Advice

The present application concerns import and processing for feed and food purposes of the genetically modified cotton line 281-24-236x3006-210-23xMON88913. Cultivation is not part of the application. Therefore, the risk assessment focuses on the accidental spillage of cottonseeds.

Cotton plants are very sensitive to temperature. The Northwestern European climate is unsuited for cotton growth. There is no indication that the introduced traits, which confer resistance to certain lepidopteran insects and tolerance to gluphosinate-ammonium and glyphosate containing herbicides, will introduce an ability to survive in the environment. Therefore, incidental spillage of cottonseeds will not lead to feral cotton populations in Northwestern Europe. COGEM considers the current GS plan sufficient for this application concerning import and processing of cotton line 281-24-236x3006-210-23xMON88913.

In view of the above, COGEM is of the opinion that the risks for man and the environment associated with import and processing of cotton line 281-24-236x3006-210-23xMON88913 are negligible. A food/feed safety assessment is carried out by other organisations. Therefore, COGEM abstains from advice on potential risks of incidental consumption.

## References

1. COGEM (2004). Marktaanvraag C/NL/04/01 voor de import van insectenresistente en herbicidentolerante katoen. COGEM advies CGM/040504-01
2. COGEM (2004)Moleculaire karakterisatie ten behoeve van import van insectresistente en herbicidentolerante katoen. COGEM advies CGM/040527-05
3. COGEM (2007). Import and processing of herbicide tolerant cotton MON88913. COGEM advise CGM/071218-01
4. Crop Protection Compendium (2007). *Gossypium* and *Gossypium hirsutum* (cotton). CD-ROM edition, © Cab International 2007, Nosworthy way, Wallingford, UK
5. OECD (2008). Consensus document on the biology of cotton (*Gossypium* spp.)
6. OGTR (2008). The biology of *Gossypium hirsutum* L. and *Gossypium barbadense* L. (cotton) Version 2 february 2008
7. European Commission (2008). Reform of the support scheme for cotton: [http://ec.europa.eu/agriculture/cap-post-2013/legal-proposals/com625/625\\_en.pdf](http://ec.europa.eu/agriculture/cap-post-2013/legal-proposals/com625/625_en.pdf) (March 8th, 2013)
8. Andersson MS & Carmen de Vicente M (2010). Gene flow between crops and their wild relatives. The John Hopkins University Press, Baltimore, Maryland, The United States of America
9. Reddy KR *et al.* (1992). Temperature effects on Pima cotton growth and development. *Agronomy Journal* 84: 237-243
10. Unruh BL & Silvertooth JC (1997). Planting and irrigation termination timing effects on the yield of Upland and Pima cotton. *Journal of Production Agriculture* 10: 74-79
11. Polunin O (2005). *Flowers of Greece and the Balkan – a field guide*. Oxford University Press. ISBN 0-19-281998-4
12. Tutin TG *et al.* (2005). *Flora Europaeae*. Volume 2. Rosaceae to Umbelliferae. Cambridge University Press. ISBN 0 521 26662 X

13. Sanahuja *et al.* (2011). *Bacillus thuringiensis*: a century of research, development and commercial applications. *Plant Biotechnol J.* 9: 283-300
14. Broderick NA *et al.* (2006). Midgut bacteria required for *Bacillus thuringiensis* insecticidal activity. *Proc. Natl. Acad. Sci. USA.* 103: 15196-15199
15. Funke T *et al.* (2006). Molecular basis for the herbicide resistance of Roundup Ready crops. *Proc. Natl. Acad. Sci. USA.* 103: 13010-13015
16. Manderscheid R & Wild A. (1986). Studies on the mechanism of inhibition by phosphinothricin of glutamine synthetase isolated from *Triticum aestivum* L. *J. Plant Physiol.* 123: 135-142
17. Strauch E *et al.* (1988). Cloning of a phosphinothricin Nacetyltransferase gene from *Streptomyces viridochromogenes* Tü494 and its expression in *Streptomyces lividans* and *Escherichia coli*. *Gene* 63: 65-74
18. Koninklijk Nederlands Meteorologisch Instituut (2013). Klimaatatlas – Langjarig gemiddelde 1981-2010. <http://www.klimaatatlas.nl/klimaatatlas.php> (February 28th, 2013)
19. COGEM (2008). Toelichting advies GA21. Brief CGM/080117-02
20. COGEM (2011). Advies m.b.t het concept van de herziene ‘Guidance on the Post-Market Environmental Monitoring (PMEM) of GM plants’ van de EFSA. COGEM advisory report CGM/110520-01
21. EFSA Panel on Genetically Modified Organisms (2011). Guidance on the Post-Market Environmental Monitoring (PMEM) of genetically modified plants. *EFSA Journal* 9:2316
22. EFSA Panel on Genetically Modified Organisms (2011). Guidance on the Post-Market Environmental Monitoring (PMEM) of genetically modified plants. *EFSA Journal* 9:2316