

Aan de staatssecretaris van
Infrastructuur en Milieu
Mevrouw S.A.M. Dijkma
Postbus 20901
2500 EX Den Haag

DATUM 12 april 2016
KENMERK CGM/160412-01
ONDERWERP Advies import en verwerking van de genetisch gemodificeerde maïslijn
1507xMON810xMIR162xNK603

Geachte mevrouw Dijkma,

Naar aanleiding van een adviesvraag betreffende een vergunningaanvraag voor de import en verwerking van genetisch gemodificeerde maïs 1507xMON810xMIR162xNK603 (EFSA/GMO/NL/2015/127), ingediend door Pioneer Hi-Bred International, Inc., deelt de COGEM u het volgende mee.

Samenvatting:

De COGEM is gevraagd om te adviseren over import en verwerking van de genetisch gemodificeerde (gg-) maïslijn 1507xMON810xMIR162xNK603. Deze lijn brengt de genen *cry1F*, *cry1Ab*, *vip3Aa20*, *cp4 epsps* en *pat* tot expressie, waardoor het gewas resistent is tegen bepaalde insecten die tot de vlinderachtigen behoren en tolerant is voor glyfosaat en glufosinaat-ammonium bevattende herbiciden.


Maïslijn 1507xMON810xMIR162xNK603 is tot stand gekomen door de vier ouderlijnen met elkaar te kruisen. De COGEM heeft eerder positief geadviseerd over alle vier de gg-ouderlijnen. Verwildering van maïsplanten is in Nederland nooit waargenomen. Daarnaast is opslag van maïs in Nederland nagenoeg uitgesloten. De moleculaire karakterisering van 1507xMON810xMIR162xNK603 is geactualiseerd en voldoet aan de criteria van de COGEM. Er zijn geen redenen om aan te nemen dat expressie van de ingebrachte genen ervoor zorgt dat de gg-maïs zou kunnen verwilderen. Bovendien zijn er in Europa geen wilde verwanten van maïs aanwezig, waardoor de ingebrachte sequenties zich niet naar andere soorten kunnen verspreiden.

Gezien het bovenstaande acht de COGEM de milieurisico's van import en verwerking van gg-maïs 1507xMON810xMIR162xNK603 verwaarloosbaar klein. Omdat andere instanties een voedselveiligheidsbeoordeling uitvoeren, heeft de COGEM bij deze vergunningaanvraag de risico's van incidentele consumptie niet beoordeeld.



De door de COGEM gehanteerde overwegingen en het hieruit voortvloeiende advies treft u hierbij aan als bijlage.

Hoogachtend,



Prof. dr. ing. Sybe Schaap
Voorzitter COGEM

c.c. Drs. H.P. de Wijs, Hoofd Bureau ggo
 Mr. J.K.B.H. Kwisthout, Ministerie van IenM

Import and processing of insect resistant and herbicide tolerant genetically modified maize 1507xMON810xMIR162xNK603

COGEM advice CGM/160412-01

- The present application (EFSA/GMO/NL/2015/127) concerns import and processing for use in feed and food of genetically modified maize 1507xMON810xMIR162xNK603;
- 1507xMON810xMIR162xNK603 was produced by conventional crossbreeding of the parental GM lines 1507, MON810, MIR162 and NK603;
- COGEM advised positively on import and processing of all four parental lines;
- The GM line expresses the *cry1F*, *cry1Ab*, *vip3Aa20*, *pmi*, *cp4 epsps* and *pat* genes;
- 1507xMON810xMIR162xNK603 is resistant to certain lepidopteran insects, tolerant to glyphosate and glufosinate-ammonium containing herbicides and has the ability to utilize mannose as the primary carbon source;
- The molecular characterization of 1507xMON810xMIR162xNK603 is updated and meets the criteria of COGEM;
- In the Netherlands feral maize populations cannot occur;
- Hybridisation of maize with other species is not possible in the Netherlands;
- The molecular characterization of 1507xMON810xMIR162xNK603 maize is adequate;
- There are no indications that expression of the introduced traits or any combination thereof will alter the fitness of 1507xMON810xMIR162xNK603 maize;
- Import and processing of maize 1507xMON810xMIR162xNK603 poses a negligible risk to the environment in the Netherlands;
- COGEM abstains from giving advice on the potential risks of incidental consumption since a food/feed assessment is already carried out by other organisations.

1. Introduction

The scope of the present application (EFSA/GMO/NL/2015/127) filed by Pioneer Hi-Bred International, Inc. is import and processing of genetically modified (GM) maize 1507xMON810xMIR162xNK603. It expresses the *cry1F*, *cry1Ab* and *vip3Aa20* genes conferring resistance to certain lepidopteran insects. In addition, the *pat* and *cp4 epsps* proteins are expressed conferring tolerance to glyphosate and glufosinate-ammonium containing herbicides. 1507xMON810xMIR162xNK603 also expresses the *pmi* gene which allows the plant to utilize mannose as the primary carbon source. The *pmi* gene is used as a selection marker.

1507xMON810xMIR162xNK603 maize was produced by conventional crossbreeding of GM maize lines 1507, MON810, MIR162 and NK603. EFSA issued positive opinions on all four parental lines for food and feed uses, import and processing.^{1,2,3,4}

The traits of the stacked GM maize will segregate in its progeny, leading to different combinations of traits being present in the kernels (seeds). Therefore, COGEM will assess the risk of 1507xMON810xMIR162xNK603 and all possible subcombinations.

Previous COGEM advices

COGEM advised positively on import and processing of parental lines 1507, MON810, MIR162 and NK603^{5,6,7,8} and on import and processing of several crosses including 1507xNK603, NK603xMON810, MON89034x1507xNK603, 1507x59122xMON810xNK603 and Bt11xMIR162x1507xGA21.^{9,10,11,12,13} COGEM also advised positively on cultivation of 1507, MON810 and NK603.^{14,15,16}

2. Environmental risk assessment

2.1 Aspects of the wild-type crop

Maize (*Zea mays*) is a member of the grass family *Poaceae*. Maize is a highly domesticated crop, originating from Central America, but nowadays maize is cultivated globally. Maize is predominantly wind pollinated.^{17,18} Insect pollination is limited since the female flowers do not produce nectar and therefore are not attractive to insect pollinators.¹⁹ In Europe, no wild relatives of maize are present and thus hybridisation with other species cannot occur.

In the Netherlands, the appearance of volunteers is very rare to absent.²⁰ Domesticated maize requires warm conditions in order to grow and does not tolerate prolonged cold and frost.^{19,21} The kernels remain on the cob after ripening and do not shatter naturally.^{19,22} In cultivation areas with warmer climatic conditions, the appearance of volunteers can occur the year following maize cultivation due to spilled cobs or kernels. However, these volunteers are usually killed by common mechanical pre-planting soil preparation practices.¹⁹

Maize is very sensitive to weed competition.²³ During the long process of domestication, maize has lost the ability to survive in the wild.¹⁸ Establishment of maize plants in the wild has never been observed in the Netherlands and COGEM is not aware of any reports of feral maize populations elsewhere in Europe.

Conclusion: In the Netherlands feral maize populations cannot occur. Hybridisation of maize with other species is not possible in the Netherlands.

2.2 Description of the introduced genes and traits

1507xMON810xMIR162xNK603 maize was produced by conventional crossbreeding of GM maize lines 1507, MON810, MIR162 and NK603. COGEM previously evaluated the molecular characterization of all parental lines and considered them to be adequate.^{5,7,8,15} The bioinformatic

analyses were updated using recent databases. COGEM is of the opinion that the molecular characterization of 1507xMON810xMIR162xNK603 is adequate.

Introduced genes	Encoded protein	Trait
<i>cry1F</i>	Cry1F δ -endotoxin originating from <i>Bacillus thuringiensis</i> subsp. <i>aizawai</i> ¹²	Lepidopteran insect resistance
<i>cry1Ab</i>	Cry1Ab δ -endotoxin originating from <i>B. thuringiensis</i> subsp. <i>kurstaki</i> ¹²	Lepidopteran insect resistance
<i>vip3Aa20</i>	Vegetative insecticidal protein (vip3Aa variant) originating from <i>B. thuringiensis</i> strain AB88 ⁷	Lepidopteran insect resistance
<i>cp4 epsps</i>	Herbicide tolerant form of 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS) enzyme originating from <i>Agrobacterium tumefaciens</i> strain CP4 ¹²	Tolerance to glyphosate containing herbicides
<i>pat</i>	Phosphinothricin N-acetyltransferase (PAT) enzyme originating from <i>Streptomyces viridochromogenes</i> strain Tü 494 ¹²	Tolerance to glufosinate-ammonium herbicides
<i>pmi</i>	Phosphomannose Isomerase (PMI) enzyme originating from <i>Escherichia coli</i> strain K-12 ⁷	Metabolizes mannose and allows for easy recovery of transformed plants
For a detailed description of the introduced genes and traits, see references.		

Conclusion: The molecular characterization of 1507xMON810xMIR162xNK603 maize is adequate. There are no indications that expression of the introduced traits or any combination thereof will alter the fitness of 1507xMON810xMIR162xNK603 maize.

2.3 Food/feed assessment

This application is submitted under Regulation (EC) 1829/2003, therefore a food/feed assessment is carried out by EFSA and national organisations involved in the assessment of food safety. In the Netherlands, a food and/or feed assessment for Regulation (EC) 1829/2003 applications is carried out by RIKILT. The outcome of the assessment by other organisations (EFSA, RIKILT) was not known when this advice was completed.

3. General Surveillance

COGEM has published several recommendations for further improvement of the general surveillance (GS) plan^{24,25} but considers the current GS plan adequate for import and processing of 1507xMON810xMIR162xNK603.

4. Overall conclusion

Import and processing of maize 1507xMON810xMIR162xNK603 including subcombinations that might be present in the kernels poses a negligible risk to the environment in the Netherlands. COGEM abstains from giving advice on the potential risks of incidental consumption since a food/feed assessment is already carried out by other organisations.

References

1. European Food Safety Authority (EFSA) (2009a). Application (EFSA-GMO-RX-1507) for renewal of authorisation for the continued marketing of existing products produced from maize 1507 for feed use, under Regulation (EC) No 1829/2003 from Pioneer Hi-Bred International, Inc./Mycogen Seeds. The EFSA Journal 1138, 1-11
2. European Food Safety Authority (EFSA) (2009b). Applications (EFSA-GMO-RX-MON810) for renewal of authorisation for the continued marketing of (1) existing food and food ingredients produced from genetically modified insect resistant maize MON810; (2) feed consisting of and/or containing maize MON810, including the use of seed for cultivation; and of (3) food and feed additives, and feed materials produced from maize MON810, all under Regulation (EC) No 1829/2003 from Monsanto. The EFSA Journal 1149, 1-85
3. European Food Safety Authority (EFSA) (2012). Scientific Opinion on application (EFSA-GMO-DE-2010-82) for the placing on the market of insect-resistant genetically modified maize MIR162 for food and feed uses, import and processing under Regulation (EC) No 1829/2003 from Syngenta. The EFSA Journal 10(6) :2756
4. Applications (references EFSA-GMO-NL-2005-22, EFSA-GMO-RX-NK603) for the placing on the market of the genetically modified glyphosate tolerant maize NK603 for cultivation, food and feed uses, import and processing and for renewal of the authorisation of maize NK603 as existing products, both under Regulation (EC) No 1829/2003 from Monsanto
5. COGEM (2015). Renewal of the authorisation for import and processing of genetically modified maize line 1507. COGEM advice CGM/150928-01
6. COGEM (1996). Markttoelating MON810. COGEM advice CGM/960807-01
7. COGEM (2010). Import and processing of genetically modified maize MIR162. COGEM advice CGM/101019-04
8. COGEM (2003). Markttoelating 'NK603 maize tolerant to glyphosate' COGEM advice CGM/030319-08
9. COGEM (2005). Import and processing of insect resistant and herbicide tolerant maize 1507 x NK603. COGEM advice CGM/050526-01
10. COGEM (2004). Marktdossier C/GB/02/M3/3 "Insectenresistente en herbicidentolerante maïs NK603 x MON810. COGEM advice CGM/040421-01
11. COGEM (2009). Import and processing of genetically modified MON89034x1507xNK603 maize. COGEM advice CGM/090930-01
12. COGEM (2012). Import and processing of genetically modified maize line 1507x59122xMON810

- xNK603. COGEM advice CGM/120402-01
13. Import and processing of genetically modified maize line Bt11xMIR162x1507xGA21
 14. COGEM (2003). Insect resistant and glufosinate ammonium tolerant transformation event 1507 maize. COGEM advice CGM/030919-04
 15. COGEM (2008). Renewal application cultivation of genetically modified maize MON810. COGEM advice CGM/080414-01
 16. COGEM (2006). Teelt van Maislijn NK603. COGEM advice CGM/060704-01
 17. Hin CJA (2001). Landbouwkundige risico's van uitkruising van GGO-gewassen. Centrum voor Landbouw en Milieu (CLM)
 18. Treu R & Emberlin J (2000). Pollen Dispersal in the Crops Maize (*Zea mays*), Oil Seed Rape (*Brassica napus* ssp. *oleifera*), Potatoes (*Solanum tuberosum*), Sugar Beet (*Beta vulgaris* ssp. *vulgaris*) and Wheat (*Triticum aestivum*). Evidence from Publications. Soil Association
 19. Andersson M & Carmen de Vicente M. (2010). Gene flow between crops and their wild relatives. The John Hopkins University Press, Baltimore, Maryland, The United States of America
 20. van de Wiel CCM et al. (2011). Crop volunteers and climate change. Effects of future climate change on the occurrence of maize, sugar beet and potato volunteers in the Netherlands. COGEM research report 2011-11
 21. Miedema P (1982). The effect of low temperature on *Zea mays*. *Advances in Agronomy* 35: 93-128
 22. Organisation for Economic Cooperation and Development (OECD) (2003). Consensus Document on the Biology of *Zea mays* subsp. *mays* (Maize)
 23. CAB International (2007). Crop Protection Compendium. *Zea mays* (maize). CD-ROM edition, Wallingford
 24. COGEM (2010). General Surveillance. COGEM report CGM/100226-01
 25. COGEM (2015). Advice on improving the general surveillance of GM crops. COGEM advice CGM/150601-02