

Aan de staatssecretaris van  
Infrastructuur en Waterstaat  
drs. V.L.W.A. Heijnen  
Postbus 20901  
2500 EX Den Haag

**DATUM** 15 april 2024  
**KENMERK** CGM/240415-01  
**ONDERWERP** Advies import en verwerking van gg-maïs DP202216xNK603xDAS-40278-9

Geachte mevrouw Heijnen,

Naar aanleiding van een vergunningaanvraag voor import en verwerking van genetisch gemodificeerde maïs DP202216xNK603xDAS-40278-9 (GMFF-2022-6232), ingediend door Corteva Agriscience LCC, deelt de COGEM u het volgende mee.

**Samenvatting:**

De COGEM is gevraagd te adviseren over de mogelijke milieurisico's van import en verwerking van de genetisch gemodificeerde (gg-)maïs DP202216xNK603xDAS-40278-9 en alle subcombinaties daarvan. In deze gg-maïs komt een gemodificeerd *pat* gen (*mopat*), een *aad-1* gen en twee kopieën van het *cp4 epsps* gen tot expressie, waardoor de plant tolerant is voor bepaalde herbiciden. Verder brengt deze gg-maïs het endogene *zmm28* gen verhoogd en langer tot expressie, waardoor de maïsplanten potentieel een hogere graanopbrengst hebben.

Verwildering van maïsplanten is in Nederland nooit waargenomen. Bovendien komt de wilde verwant van maïs, teosinte, niet in de natuurlijke omgeving in Nederland voor, waardoor de ingebrachte sequenties zich niet naar andere soorten kunnen verspreiden.

De moleculaire karakterisering van DP202216xNK603xDAS-40278-9 voldoet aan de eisen van de COGEM. Er zijn geen redenen om aan te nemen dat expressie van de ingebrachte genen ervoor zorgt dat deze gg-maïs zou kunnen verwilderen.

Gezien het bovenstaande acht de COGEM de milieurisico's van de import en verwerking van de gg-maïs DP202216xNK603xDAS-40278-9, en sub-combinaties hiervan, verwaarloosbaar klein.

Omdat een voedselveiligheidsbeoordeling door andere instanties wordt uitgevoerd, heeft de COGEM bij deze vergunningaanvraag de risico's van incidentele consumptie niet beoordeeld.



De door de COGEM gehanteerde overwegingen en het hieruit voortvloeiende advies treft u hierbij aan als bijlage.

Hoogachtend,

Prof. dr. ing. Sybe Schaap  
Voorzitter COGEM

- c.c.
- Drs. Y de Keulenaar, Hoofd Bureau ggo
  - Ministerie van IenW, Directie Omgevingsveiligheid en milieurisico's,  
DG Milieu en Internationaal
  - Ing. M.A.C. Möllers, Food-Feed loket

## Import and processing of genetically modified maize DP202216xNK603xDAS-40278-9 and subcombinations

### COGEM advice CGM/240415-01

- The present application (GMFF-2022-6232) concerns the authorisation of genetically modified (GM) maize DP202216xNK603xDAS-40278-9 and its subcombinations for import and processing for use in feed and food;
- The stacked event maize DP202216xNK603xDAS-40278-9 was produced by conventional crossbreeding of the three GM parental maize lines;
- Previously, COGEM advised positively on the import and processing of all three parental lines;
- The GM maize expresses a modified *pat* gene (*mo-pat*), the *aad-1* gene and two copies of the *cp4 epsps* gene, conferring tolerance to glyphosate, glufosinate-ammonium, 2,4 dichlorophenoxyacetic acid (2,4-D), and aryloxyphenoxypropionate (AOPP) containing herbicides. Furthermore, it expresses the *zmm28* gene leading to increased and extended production of the endogenous ZMM28 protein resulting in plants with an enhanced grain yield potential;
- In the Netherlands, feral maize populations have never been observed and the appearance of volunteers is rare;
- In the Netherlands, the wild relative of maize, teosinte is not present in the environment, hybridisation of GM maize with other species is therefore not possible;
- The molecular characterisation of DP202216xNK603xDAS-40278-9 has been updated and meets the criteria of COGEM;
- The updated bioinformatic analysis did not provide indications of potential environmental risks;
- There are no indications that the introduced traits allow the stacked event maize DP202216xNK603xDAS-40278-9 to survive in the Netherlands;
- There are no indications that DP202216xNK603xDAS-40278-9 could establish feral populations in the Netherlands;
- COGEM is of the opinion that import and processing of maize DP202216xNK603xDAS-40278-9, and GM maize consisting of subcombinations of its parental GM maize lines poses a negligible risk to the environment in the Netherlands;
- COGEM abstains from giving advice on the potential risks of incidental consumption since a food/feed assessment is carried out by other organisations.

## 1. Introduction

The present application (GMFF-2022-6232) filed by Corteva Agriscience LCC concerns import and processing of DP202216xNK603xDAS-40278-9 (DP-202216-6xMON-00603-6xDAS-40278-9) maize and genetically modified (GM) maize consisting of its subcombinations. The GM maize was produced by conventional crossbreeding of the three genetically modified (GM) parental maize lines. It expresses the *mo-pat* and *aad-1* genes, and two copies of the *cp4 epsps* gene, conferring tolerance to glufosinate-ammonium, glyphosate, 2,4-D and AOPP containing herbicides. Furthermore, it expresses the *zmm28* gene leading to increased and extended production of the endogenous ZMM28 protein resulting in plants with an enhanced grain yield potential.<sup>1,2</sup>

The parental lines NK603<sup>3</sup> and DAS-40278-9<sup>4</sup>, and the NK603xDAS-40278-9<sup>5</sup> subcombination have been authorised for import and processing for use in food and feed in the European Union. The import, food and feed of the parental line DP202216 is not authorized yet.

## 2. Previous COGEM advice

Previously COGEM advised positively on cultivation of the parental line NK603<sup>6,7</sup>, and on import and processing of the parental lines DP202216<sup>8</sup> and DAS-40278-9<sup>9</sup>. COGEM also advised positively on the import and processing of several stacked events including MON89034x1507xNK603xDAS-40278-9<sup>10</sup> and MON89034x1507xMIR162x NK603xDAS-40278-9<sup>11</sup>.

## 3. Environmental risk assessment

### 3.1 Characteristics of maize

Maize (*Zea mays*) is a member of the grass family *Poaceae*. It is a highly domesticated crop originating from Central America, but nowadays cultivated globally. Maize is wind pollinated,<sup>12,13</sup> and has both male and female flowers that are spatially separated. The female flowers are not attractive to insect pollinators, because they do not produce nectar. Insect pollination of maize is highly limited but cannot be excluded.<sup>14</sup>

In Europe, no wild relatives of maize other than teosinte, are present and, therefore, hybridization of GM maize with other species cannot occur. Teosinte has been detected in Spain<sup>15,16,17</sup> and France.<sup>18,19</sup> In these countries, it is present as a weed in some arable fields. In the Netherlands, teosinte is not present in the environment.<sup>20</sup>

Maize requires warm conditions to grow.<sup>14,21</sup> In cultivation areas with warm climatic conditions, the appearance of volunteers can occur the year following maize cultivation due to spilled cobs or kernels. However, these volunteers are usually killed by common mechanical pre-planting soil preparation practices.<sup>14</sup> Maize does not tolerate prolonged cold and frost.

Maize is very sensitive to weed competition.<sup>22</sup> During the long process of domestication, maize has lost the ability to persist in the wild.<sup>13</sup> A soil seed bank, small seeds, and an extended period of flowering and seed production are characteristics often observed in persistent weeds.<sup>23</sup> Maize lacks all these characteristics. After ripening, the seeds (the kernels) adhere to the cob and do not scatter naturally.<sup>14,24</sup> Consequently, seed dispersal is severely hampered.

During field observations in Austria some volunteers and maize plants were observed in non-agricultural habitats.<sup>25</sup> In the Netherlands, the appearance of volunteers is rare, although maize plants occasionally have been observed outside agricultural fields.<sup>20,26</sup> Any volunteers emerging will be killed by frost at the onset of winter.<sup>20</sup> COGEM is not aware of any reports of feral maize populations in the Netherlands.<sup>20</sup>

**Conclusion:** In the Netherlands, feral maize populations do not occur and, therefore, hybridisation of maize with other species is impossible.

### 3.2 Description of the introduced genes, traits, and regulatory elements

Maize DP202216xNK603xDAS-40278-9 was created by conventional cross-breeding of the parental lines. For a description of the parental lines, see previous COGEM advices.<sup>6,7,8,9,10,11</sup> A description of the inserted genetic elements is listed in the table below. The list is limited to information on the introduced genes, corresponding traits, and regulatory elements (promoters and terminators).

Introduced genes	Encoded proteins	Traits	Regulatory elements
<i>aad-1</i> (DAS-40278-9)	Aryloxyalkanoate dioxygenase (AAD-1) enzyme originating from <i>Sphingobium herbicidovorans</i> <sup>27</sup>	Cleavage of several synthetic auxins and certain 'aryloxyphenoxy-propionate' (AOPP) herbicides, resulting in a tolerance to several synthetic auxin acting herbicides like 2,4 dichlorophenoxyacetic acid' (2,4-D) and to AOPP containing herbicides	Ubiquitin promoter (ZmUbi1) and ZmPer5 3'UTR from <i>Zea mays</i>
<i>cp4 epsps</i> (NK603)	5-enolpyruvylshikimate-3-phosphate synthase (EPSPS) from <i>Agrobacterium</i> sp. strain CP4 <sup>28,29</sup>	Tolerance to glyphosate containing herbicides	Rice actin ( <i>p-ract1</i> ) promoter and intron from <i>Oryza sativa</i> , <i>ctp2</i> transit peptide from ( <i>A. thaliana</i> ) and nopaline synthase ( <i>NOS</i> 3'UTR) terminator from <i>A. tumefaciens</i>
<i>cp4 epsps</i> <i>L214P</i> (NK603)	5-enolpyruvylshikimate-3-phosphate synthase from <i>Agrobacterium</i> sp. strain CP4 <sup>28,29</sup>	Tolerance to glyphosate containing herbicides	Enhanced 35S ( <i>e35S</i> ) promoter from <i>Cauliflower mosaic virus</i> (CaMV),

			<i>Zmhsp70</i> intron ( <i>Zea mays</i> ), <i>ctp2</i> transit peptide from ( <i>A. thaliana</i> ) nopaline synthase ( <i>NOS</i> 3'UTR) terminator from <i>A. tumefaciens</i>
<i>zmm28</i> (DP202216)	MADS-domain transcription factor gene originating from <i>Zea mays</i> encoding for ZMM28 protein <sup>2,1,30,31</sup>	Plants with enhanced grain yield potential	Constitutive promoter ( <i>zm-gos2</i> ) and intron from the ubiquitin gene 1 ( <i>ubiZM1</i> ) from <i>Z. mays</i> and proteinase inhibitor II ( <i>pinII</i> ) gene terminator from <i>Solanum tuberosum</i>
<i>mo-pat</i> (DP202216)	Maize codon-optimised version of phosphinothricin acetyl transferase ( <i>pat</i> ) gene from <i>Streptomyces viridochromogenes</i> <sup>32,33,34</sup>	Tolerance to glufosinate-ammonium containing herbicides	Promoter, 5'UTR and intron from the <i>ubiZM1</i> gene from <i>Z. mays</i> and <i>pinII</i> terminator from <i>S. tuberosum</i>
See references for a detailed description of the traits			

### 3.3 Molecular characterisation

Previously, COGEM evaluated the molecular characterisation of each parental line and considered these to be adequate.<sup>6,7,8,9,10,11</sup>

The applicant sequenced the DP202216xNK603xDAS-40278-9 maize and compared the obtained DP202216, NK603 and DAS-40278-9 inserts and flanking sequences in DP202216xNK603xDAS-40278-9 with the sequences of the corresponding single events. According to the applicant the DP202216, NK603 and DAS-40278-9 inserts and flanking DNA regions in the stacked event maize are identical to the DNA sequences determined previously for the respective single events.

The applicant updated the bioinformatics analyses of the DP202216, NK603 and DAS-40278-9 inserts and of the 5' and 3' flanking regions using recent databases. According to the applicant, the putative products of the open reading frames spanning the 5' and 3' junctions of the inserts did not generate any protein sequence similarity with known allergens, toxins, or other biologically active proteins.

The molecular characterisation was conducted according to the criteria previously laid down by COGEM.<sup>35</sup> The results from the updated molecular characterisation do not provide indications that the stacked event maize DP202216xNK603xDAS-40278-9 could pose a risk to the environment.

**Conclusion:** The molecular characterisation of maize DP202216xNK603xDAS-40278-9 is adequate and no indications of potential environmental risks were identified.

### 3.4 Phenotypic and agronomic characteristics

Previously, COGEM evaluated the phenotypic and agronomic characteristics of each parental line of DP202216xNK603xDAS-40278-9, and found no deviations influencing the outcome of the environmental risk assessment.

The applicant analysed the phenotypic and agronomic characteristics of DP202216xNK603xDAS-40278-9 and noted that these, except for the introduced traits, are not different to the non-GM near-isogenic control line, and are equivalent to the reference varieties, taking into account natural variation. The results of the phenotypic evaluation do not give reason to assume that the GM maize could pose an environmental risk.

Therefore, COGEM is of the opinion that there are no indications that the introduced traits allow maize DP202216xNK603xDAS-40278-9 to survive or establish in the Dutch environment.

**Conclusion:** There are no indications that the introduced traits allow DP202216xNK603xDAS-40278-9 to survive in the Netherlands. DP202216xNK603xDAS-40278-9 has no increased potential for the establishment of feral populations in the Netherlands.

#### **4. Food/feed assessment**

This application is submitted under Regulation (EC) 1829/2003, therefore a food/feed assessment is carried out by EFSA and national organisations involved in the assessment of food safety. In the Netherlands, a food and/or feed assessment for Regulation (EC) 1829/2003 applications is carried out by Wageningen Food Safety Research (WFSR). The outcome of the assessment by other organisations (EFSA, WFSR) was not known when this advice was completed.

#### **5. Post-market environmental monitoring (PMEM)**

The applicant supplied a general surveillance plan as part of the PMEM. COGEM has published several recommendations for further improvement of the general surveillance (GS) plan,<sup>36,37</sup> but considers the current GS plan adequate for the import and processing of maize DP202216xNK603xDAS-40278-9.

#### **6. Overall conclusion**

COGEM is of the opinion that import and processing of DP202216xNK603xDAS-40278-9 maize and GM maize consisting of subcombinations of its parental GM maize lines pose a negligible risk to the environment in the Netherlands. COGEM abstains from giving advice on the potential risks of incidental consumption since other organisations carry out a food/feed assessment.

#### **References**

1. Anderson JA *et al.* (2019). Evaluation of the history of safe use of the maize ZMM28 protein. *J. Agric. Food Chem.* 67: 7466-7474

2. Wu J *et al.* (2019). Overexpression of zmm28 increases maize grain yield in the field. *Proc. Natl. Acad. Sci. U.S.A.* doi: 10.1073/pnas.1902593116
3. European Commission (2015). Commission Implementing Decision (EU) 2015/684 of 24 April 2015 authorising the placing on the market of genetically modified maize NK603 (MON-ØØ6Ø3-6) and renewing the existing maize NK603 (MON-ØØ6Ø3-6) products, pursuant to Regulation (EC) No 1829/2003 of the European Parliament and of the Council. (C(2015) 2753). *Official Journal of the European Union*. L 112/6
4. European Commission (2017). Commission Implementing Decision authorising the placing on the market of products containing, consisting of, or produced from genetically modified maize DAS-40278-9, pursuant to Regulation (EC) No 1829/2003 of the European parliament and of the Council on genetically modified food and feed (2017/1212). *Official Journal of the European Union* 6.7.2017 L173/43
5. European Commission (2017). Commission Implementing Decision authorising the placing on the market of products containing, consisting of or produced from genetically modified maize MON 89034 × 1507 × MIR162 × NK603 × DAS-40278-9 and nine sub-combinations, pursuant to Regulation (EC) No 1829/2003 of the European Parliament and of the Council on genetically modified food and feed (C(2023) 2133). *Official Journal of the European Union*.
6. COGEM (2003). Markttoelating 'NK603 maize tolerant for glyphosate'. COGEM advice CGM/030319-08 [In Dutch]
7. COGEM (2006). Cultivation of herbicide tolerant maize line NK603. COGEM advice CGM/060704-01
8. COGEM (2019). Import and processing of herbicide tolerant genetically modified maize DP202216 with enhanced grain yield potential COGEM advice CGM/191129-04
9. COGEM (2011). Import and processing of genetically modified maize DAS-40728-9. COGEM advice CGM/110510-01
10. COGEM (2017). Import and processing of genetically modified maize MON89034x1507xNK603xDAS-40278-9. COGEM advice CGM/170131-01
11. COGEM (2018). Import and processing of genetically modified maize MON89034x1507xMIR162xNK603xDAS-40278-9 and sub-combinations. COGEM advice CGM/181220-03
12. Hin CJA (2001). Landbouwkundige risico's van uitkruising van GGO-gewassen. Centrum voor Landbouw en Milieu (CLM) [In Dutch]
13. Treu R & Emberlin J (2000). Pollen dispersal in the crops Maize (*Zea mays*), Oil Seed Rape (*Brassica napus* ssp. *oleifera*), Potatoes (*Solanum tuberosum*), Sugar Beet (*Beta vulgaris* ssp. *vulgaris*) and Wheat (*Triticum aestivum*). Evidence from Publications. Soil Association
14. Andersson M & Carmen de Vicente M (2010). Gene flow between crops and their wild relatives. The John Hopkins University Press, Baltimore, Maryland, The United States of America
15. Pardo G *et al.* (2014). El Teosinte (*Zea mays*, spp.). Cent. Sanid. y Certificación Veg. DGA [in Spanish]
16. Pardo G *et al.* (2015). Presencia de teosinte (*Zea* spp.) como mala hierba en los regadíos del valle del Ebro. XV Congr. la Soc. Española Malherbología La Malherbología y la Transf. tecnológica Sevilla, 417-424 [In Spanish]



17. Pardo G *et al.* (2016). El teosinte: descripción, situación actual en el valle del Ebro y resultados de los primeros ensayos. *Vida Rural*.408: 42-47 [In Spanish]
18. Agri79 (2013). Téosinte - La téosinte exige une vigilance toute particulière. <https://www.agri79.fr/la-teosinte-exige-une-vigilance-toute-particuliere> (visited: March 7th 2024) [In French]
19. Trtikova M *et al.* (2017). Teosinte in Europe - Searching for the origin of a novel weed. *Sci. Rep.* 7: 1560. doi: 10.1038/s41598-017-01478-w
20. Huiting HF *et al.* (2018). Are teosinte and feral maize present in the Netherlands? COGEM report CGM 2018-06
21. Miedema P (1982). The effect of low temperature on *Zea mays*. *Advances in Agronomy* 35: 93-128
22. CAB International (2007). *Crop Protection Compendium. Zea mays (maize)*. CD-ROM edition, Wallingford
23. Kos SP *et al.* (2012). Can transgenic crops go wild? A literature study on using plant traits for weediness pre-screening. COGEM research report CGM 2012-01
24. Organisation for Economic Cooperation and Development (OECD) (2003). Consensus document on the biology of *Zea mays ssp. mays* (Maize)
25. Pascher K (2016). Spread of volunteer and feral maize plants in Central Europe: recent data from Austria. *Environ. Sci. Eur.* 28: 30
26. van de Wiel CCM *et al.* (2011). Crop volunteers and climate change. Effects of future climate change on the occurrence of Maize, Sugar Beet and Potato volunteers in the Netherlands. COGEM research report 2011-11
27. Herman RA *et al.* (2010). Compositional safety of event DAS-40278-9 (AAD-1) herbicide-tolerant maize. *GM Crops*. 1: 294-311
28. Funke T *et al.* (2006). Molecular basis for the herbicide resistance of Roundup Ready crops. *PNAS* 103: 13010-13015
29. Harrison *et al.* (1993). Characterization of Microbially-Expressed Protein: CP4 EPSPS. Monsanto Company, pp.
30. Münster T *et al.* (2002). Maize MADS-Box Genes Galore. *Maydica* 47: 287-301
31. Pařenicová L *et al.* (2003). Molecular and phylogenetic analyses of the complete MADS-box transcription factor family in Arabidopsis: new openings to the MADS world. *Plant Cell*. 15: 1538-1551
31. Wohlleben W *et al.* (1988). Nucleotide sequence of the phosphinothricin N-acetyltransferase gene form *Streptomyces viridochromogenes* Tü494 and its expression in *Nicotiana tabacum*. *Gene* 70: 25-37
32. Eckes P, Uijtewaal B and Donn G, 1989. A synthetic gene confers resistance to the broad spectrum herbicide L-phosphinothricin in plants. *Journal of Cellular Biochemistry Supplement* 13D, 334, Abstract M516
33. Organisation for Economic Cooperation and Development (OECD) (1999). Consensus document on general information concerning the genes and their enzymes that confer tolerance to phosphinothricin herbicide.
35. COGEM (2014). Reconsideration of the molecular characterisation criteria for marketing authorisation of GM crops. COGEM topic report CGM/140929-02
36. COGEM (2010). General Surveillance. COGEM topic report CGM/100226-01
37. COGEM (2015). Advice on improving the general surveillance of GM crops. COGEM advice CGM/150601-02