

Aan de staatssecretaris van
Infrastructuur en Waterstaat
Mr. drs. A.W.H. Bertram
Postbus 20901
2500 EX Den Haag

DATUM 02 april 2026
KENMERK CGM/260402-01
ONDERWERP Advies Hernieuwing markttoelating gg-anjer Moonvista FLO-40685-2

Geachte mevrouw Bertram,

Naar aanleiding van een adviesvraag betreffende de hernieuwing van de vergunning voor import, distributie en verkoop van snijbloemen van de gg-anjervariëteit FLO-40685-2 (Florigene® Moonvista™) van Suntory Flowers Ltd (C/NL/13/02_001), deelt de COGEM u het volgende mee. Op verzoek van het Food-Feed loket voor ggo-markttoelatingen is het advies in het Engels geschreven.

Samenvatting:


De COGEM is gevraagd te adviseren over de hernieuwing van de vergunning voor import, distributie en verkoop van snijbloemen van de genetisch gemodificeerde (gg-)anjervariëteit FLO-40685-2 (Florigene®Moonvista™) in Europa. Anjers vormen van nature geen blauw pigment. Door toevoeging van twee genen uit petunia en viooltje kan de gg-anjer dit pigment wel maken en krijgt de bloem een 'blauwe' kleur. De eerste vergunning is in 2019 voor een periode van tien jaar afgegeven. De onderhavige aanvraag betreft een verlenging van deze vergunning.

De cultuur-anjer is niet in staat tot verwildering. De ingebrachte genen hebben geen invloed op het verwilderingspotentieel. De kans op verspreiding van genen van de gg-anjer naar verwante soorten in de natuur is verwaarloosbaar klein. In de geactualiseerde bioinformatische analyses, literatuurstudie, en monitoringsrapporten van de afgelopen jaren zijn geen mogelijke risico's voor mens en milieu geïdentificeerd.

De gg-anjer mag niet worden gebruikt voor voedsel doeleinden. Soms worden bloemblaadjes van anjer gebruikt als garnering. Er is geen reden om aan te nemen dat incidentele consumptie risico's met zich meebrengt. Concluderend is de COGEM van oordeel dat de risico's van verlenging van import, distributie en verkoop van snijbloemen van gg-anjer FLO-40685-2 voor mens en milieu verwaarloosbaar klein zijn.

De door de COGEM gehanteerde overwegingen en het hieruit voortvloeiende advies treft u hierbij aan als bijlage.

Hoogachtend,



Prof. dr. ing. Sybe Schaap
Voorzitter COGEM

c.c.

- Drs. Y. de Keulenaar, Hoofd Bureau ggo
- Ministerie van IenW, Directie Omgevingsveiligheid en milieurisico's, DG Milieu en Internationaal
- Ing. M.A.C. Möllers, Food-Feed loket

Renewal of the authorisation for import, distribution and retail of genetically modified carnation FLO-40685-2 (Moonvista™)

COGEM advice CGM/260402-01

1. Introduction

COGEM has been requested to evaluate the environmental risks associated with the renewal of the authorisation for import, distribution and retail of cut flowers of genetically modified (GM) carnation (*Dianthus caryophyllus*) FLO-40685-2 (also known as Florigene®Moonvista™). The present application (C/NL/13/02_001), was filed by Suntory Flowers Ltd.

Consent to place Moonvista™ on the market in Europe was granted in 2019.¹ Since import authorisations remain valid for a period of 10 years, the applicant filed an application for a renewal of the authorisation. COGEM previously advised positively on the import, distribution and retail in the EU of GM carnation Moonvista™, FLO-40685-2 in 2013.² COGEM has also issued positive opinions on import, distribution and retail of similar GM carnations with altered flower colours (e.g., Moonvelvet™, Moonlite™, Moonberry™ and Moonaqua™).^{3,4,5,6,7,8,9,10,11,12,13}

1.1 Carnation

Carnation belongs to the species *Dianthus caryophyllus* of the widely cultivated genus *Dianthus*. Wild *Dianthus* species are found throughout Europe, the Balkan and the Mediterranean area.^{14,15,16} In the Netherlands, several native *Dianthus* species occur and some *Dianthus* species that are commonly grown as garden plants have established themselves in the wild.^{17,18}

Carnations have been cultured for hundreds of years to improve flower size and generate colour variants. Cultivated carnations are almost certainly hybrids between two or more *Dianthus* species, one of which is most likely *D. caryophyllus*. Cultivated carnation is highly domesticated; it is not propagated by seed, but vegetatively by cuttings and tissue culture. Carnation also does not spread vegetatively spontaneously, or produce vegetative organs like bulbs, stolons or rhizomes.¹⁹ Carnation is semi-winter hardy, has no weedy characteristics and even after decades of cultivation has never shown to be able to establish itself in the wild.²⁰

In nature, pollination of *D. caryophyllus* occurs exclusively by lepidopteran insects. *Dianthus* carnation pollen cannot be spread by wind. Any pollen produced is heavy, sticky and deeply buried in the flower.^{21,22} Although the Netherlands has a large carnation handling industry, carnation pollen is not detected in the atmosphere.^{23,24,25} Breeding has increased the number of petals present in carnation cultivars. As a result, the reproductive tissues of the flower have become enclosed, restricting access to insect pollinators.¹⁴ There has never been any evidence of spontaneous hybridisation between carnation and wild *Dianthus* species, despite the fact that carnation has been cultivated worldwide for centuries.^{14,26}

1.2 GM carnation FLO-40685-2

GM carnation FLO-40685-2 was developed by *Agrobacterium*-mediated transformation, and expresses the genes *f3'5'h* and *dfr* which result in an altered flower colour, and *SuRB* which provides tolerance to

sulfonylurea herbicides and is used as marker trait in the selection of GM plants. For an overview and description of the introduced genes and traits in FLO-40685-2, see Table 1.

Table 1. Description of the introduced genes and traits.

Introduced genes	Encoded proteins	Regulatory elements	Traits
SuRB	Acetolactate synthase (ALS) enzyme derived from <i>Nicotiana tabacum</i> . ²⁷	Cauliflower mosaic virus (CaMV) 35S promoter, 5' untranslated region from the cDNA corresponding to the petunia gene encoding chlorophyll a/b binding protein. Terminator from <i>N. tabacum</i> SuRB gene. ^{2,28}	Confers tolerance to ALS inhibiting (sulfonylurea) herbicides which are used to select transformants.
<i>f3'5'h</i>	Flavonoid 3'5' hydroxylase (F3'5'H) enzyme derived from <i>Viola hortensis</i> . ²⁹	Promoter from snapdragon (<i>Antirrhinum majus</i>) chalcone synthase (CHS) gene, and terminator sequence from the petunia D8 gene encoding a phospholipid transfer protein homologue. ^{2,28}	Modified flower colour.
<i>dfr</i>	Dihydroflavonol-4-reductase (DFR) derived from <i>Petunia × hybrida</i> . ³⁰	Own promoter and terminator of the petunia <i>dfr</i> gene. ^{2,28}	Modified flower colour.

2. Environmental risk assessment

The application for market authorisation of GM carnation FLO-40685-2 for import, distribution and retail of cut flowers contains reports of the annual monitoring reports, bioinformatic analyses, and other new information relating to FLO-40685-2 that has become available since the consent to market was issued. Integration of the T-DNA (fully or partially) in FLO-40685-2 has occurred at four loci. All four transgenic loci were re-sequenced after submission of the marketing application. In locus 1, a three base pair correction was required, whereas sequences of loci 2, 3, and 4 were correct in the original application. Open reading frames (ORFs) were generated from the updated four loci sequences of FLO-40685-2. Using the translated protein sequences of ORFs bridging all T-DNA-genomic DNA junctions, bio-informatic analyses were performed to assess potential protein sequence homology to allergens and toxins with up-to-date databases. In addition, translated protein sequences of the T-DNA coding regions of the transformation vector (pCGP1991) were assessed for potential homology to toxin-related proteins or allergens. No indications for potential environmental risks were identified.

The applicant supplied annual monitoring reports for FLO-40685-2 between July 2019 and June 2025. As part of yearly monitoring, the applicant obtains questionnaire feedback from the importer and expert monitoring groups, to obtain information on illegal propagation of GM carnation in Europe. Furthermore, they contacted individual scientists and several institutions (herbaria, plant protection services, conservation groups etc.) to request any reports on the identification of wild populations of carnation, *D. caryophyllus* or related *Dianthus* species. The monitoring plan was adapted once in November 2021, discontinuing the need for the so called 'mail out' component of the monitoring plan. COGEM previously advised positively on the discontinuation of the 'mail out' component, as the alternatives methods were deemed sufficient for timely observation.³¹ This change was implemented in 2023. The applicant also inspected the composting areas of the production sites in Colombia and Ecuador and investigated whether GM carnation established at these areas. No evidence of the

establishment of FLO-40685-2 nor introgression of the transgenes into wild *Dianthus* species was obtained.

In addition, a literature and database search was performed as part of the monitoring process for FLO-40685-2, to identify reports of carnation in the wild. The literature and records evaluated by the applicant also did not contain any reports of carnations occurring in the wild or evidence of introgression of the transgenes to wild *Dianthus* species.

The genes introduced in FLO-40685-2 do not encode toxic or allergenic products. The authorisation of FLO-40685-2 is limited to import, distribution and retail and does not allow the use of the GM carnation as food. Therefore, retailers will not be allowed to sell the petals of the GM carnation for food purposes. Petals of carnation can sometimes be used in dishes and as garnishing.^{32,33,34} Occasional consumption of flower petals is unlikely to lead to an adverse effect.

COGEM is not aware of any reports indicating that incidental consumption of FLO-40685-2 poses a risk to human health. In view of the above-mentioned considerations, COGEM is of the opinion that unauthorised incidental consumption of GM carnation FLO-40685-2 poses a negligible risk to human health.

3. Conclusion

No evidence was found to indicate a change in the outcome of the risk assessment carried out at the time of the marketing approval of FLO-40685-2. COGEM is of the opinion that renewal of the authorisation for import, distribution and retail of cut flowers of GM carnation FLO-40685-2 (Moonvista™) poses a negligible risk to human health and the European environment.

References

1. Commission Implementing Decision (EU) 2019/1300 of 26 July 2019 as regards the placing on the market of a genetically modified carnation (*Dianthus caryophyllus* L., line FLO-40685-2). Official Journal of the European Union L 204/46, 2.8.2019
2. COGEM (2013). Import, distribution and retail of GM carnation FLO-40685-1. COGEM advice CGM/131217-01
3. COGEM (2007). Import of genetically modified carnation 'Moonaqua'. COGEM advice CGM/070206-02
4. COGEM (2018). Renewal of the authorisation for import, distribution and retail of genetically modified carnation FLO-40689-6. COGEM advice CGM/180613-02
5. COGEM (2009). Import, distribution and retail of GM-carnation IFD-25958-3. COGEM advice CGM/090429-01
6. COGEM (2005). Import of cut flowers of the genetically modified carnation 'Florigene Moonlite' (C/NL/04/02). COGEM advice CGM/050207-01
7. COGEM (2009). Import, distribution and retail of gm-carnation 'Moonshadow'. COGEM advice CGM/090407-08
8. COGEM (2009). Import, distribution and retail of GM-carnation IFD-25958-3. COGEM advice CGM/090429-01
9. COGEM (2009). Import, distribution and retail of GM carnation IFD-26407-2. COGEM advice CGM/090504-06
10. COGEM (2016). Renewal of the consent for import, distribution and retail of genetically modified carnation FLO-40644-6. COGEM advice CGM/160706-03
11. COGEM (2013). Import, distribution and retail of cut flowers with modified flower colour (GM carnation SHD-27531-4). COGEM advice CGM/130711-01
12. COGEM (2024). Renewal of the authorisation for import, distribution and retail of genetically modified carnation IFD-25958-3 (Moonberry). COGEM advice CGM/240502-01
13. COGEM (2024). Renewal of the authorisation for import, distribution and retail of genetically modified carnation IFD-26407-2 (Moonvelvet). COGEM advice CGM/240502-02
14. Office of the Gene Technology Regulator (2015). The biology of *Dianthus caryophyllus* L. (Carnation). Australian Government. Department of Health
15. Tela Botanica - *Dianthus caryophyllus* L. <http://www.tela-botanica.org/bdtfx-nn-22083-synthese> (visited March 10th, 2026)

16. Flora Italiana - Dianthus caryophyllus L. <http://luirig.altervista.org/flora/taxa/index1.php?scientific-name=dianthus+caryophyllus> (visited March 10th, 2026)
17. Duistermaat H. (2020). Heukels' Flora van Nederland, ed. 24. Noordhoff Uitgevers.
18. FLORON verspreidingsatlas. <https://www.verspreidingsatlas.nl/0401> (visited March 10th, 2026)
19. Strid A & Tan K (1997). Flora Hellenica, Volume 1. Koeltz scientific books. Koningstein, Germany
20. CABI (2007). Crop Protection Compendium, 2007 Edition. Wallingford, United Kingdom
21. Jennersten O (1983). Butterfly visitors as vectors of Ustilago violacea spores between Caryophyllaceae plants. Oikos 40:125-130
22. Keane AT (1989). Breeding new carnation cultivars. IPPS 39: 88-89
23. Driessen MNBM & Derksen JWM (1988). Pollenatlas van de Nederlandse Atmosfeer, Onkenhout, Hilversum, eerste druk
24. Elkerliek ziekenhuis. Pollentellingen 2026. <https://www.elkerliek.nl/elkerliek.nl/specialismen/AKL/Pollentellingen.pdf> (visited March 10th, 2026)
25. Leids Universitair Medisch Centrum. Pollentelling <https://sec.lumc.nl/pollenwebextern/> (visited March 10th, 2026)
26. Buell KM (1952). Developmental morphology in Dianthus I. Structure of the pistil and seed development. American Journal of Botany 39: 194-210
27. Keeler et al. (1993). Regulation of tobacco acetolactate synthase gene expression. Plant Physiol. 102:1009-1018
28. European Food Safety Authority (2018). Risk assessment of new sequencing information on genetically modified carnation FLO-40685-2. EFSA Journal 16: 5396
29. Fukui Y et al. (2003). A rationale for the shift in colour towards blue in transgenic carnation flowers expressing the flavonoid 3',5'-hydroxylase gene. Phytochemistry 63: 15-23
30. Beld M et al. (1989). Flavonoid synthesis in Petunia hybrid: partial characterization of duhydroflavonol-4-reductase genes. Plant Mol. Biol. 13: 491-502
31. COGEM (2021). Advies 'wijziging van het monitoringsplan voor gg-anjersdow'. COGEM advice CGM/211005-01
32. Eat me – Anjer. <https://www.eatme.nl/producten/anjer> (visited March 10th, 2026)
33. Jardins de pareillas – art de vivre entre botanique, gastronomie et créativité. <http://jardinsdepareillas.over-blog.com/article-2893237.html> (visited March 10th, 2026)
34. Elle à table. <http://www.elle.fr/Elle-a-Table/Recettes-de-cuisine/Tartines-au-beurre-d-oeillet-551895> (visited March 10th, 2026)